

## Feeding Instructions:

### Layer Birds for Eggs

Age	Product ID	Product	Average Feed Intake
0 - 42 days	110021	21% Poultry Starter Medicated Crumbles	35 g/day
0 - 42 days	110041	21% Poultry Starter Crumbles	35 g/day
42 - 126 days/1 <sup>st</sup> egg	120081	16% Poultry Grower/Finisher Crumbles	70 g/day
Laying	130061	17% Complete Poultry Layer Short Pellets	110 g/day

### Broiler Birds for Meat

Age	Product ID	Product	Average Feed Intake
0 - 28 days	110021	21% Poultry Starter Medicated Crumbles	75 g/day
0 - 28 days	110041	21% Poultry Starter Crumbles	75 g/day
28 days - market	120081	16% Poultry Grower/Finisher Crumbles	150 g/day

Always provide ample clean, fresh water. For further information refer to our poultry guide ([link poultry guide](#)).

### Warning:

Do not feed to laying birds in production.

### Caution:

- Directions for use must be carefully followed.
- Do not use Amprolium medicated feed as treatment for outbreaks of Coccidiosis.
- If unexpected deaths occur, obtain an accurate diagnosis and follow a veterinarian's or poultry pathologist's recommendations.

### Ingredients:

A list of the ingredients used in this feed may be obtained from the manufacturer or registrant.

This feed contains added **Selenium at 0.3 mg/kg**.

## Feed Analysis

Crude Protein (min) 21.0%

Phosphorus (act) 0.7%

Crude Fat (min) 2.0%

Vitamin A (min) 10,000 IU/kg

Crude Fibre (max) 6.0%

Vitamin D (min) 4,000 IU/kg

Sodium (act) 0.15%

Vitamin E (min) 50 IU/kg

Calcium (act) 1.0%

Individual results from the use of this product may vary due to management, environment, genetics, health and sanitation differences. Therefore, Trouw Nutrition Canada, Inc. does not warrant or guarantee individual results.